

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 November 2002 (14.11.2002)

PCT

(10) International Publication Number
WO 02/090324 A1

(51) International Patent Classification⁷: C07C 303/00

Kee-Young [KR/KR]; Inheon Apt. #101-602, 32-3,
Bongcheon-Dong, Kwanak-Gu, Seoul 151-050 (KR).

(21) International Application Number: PCT/KR02/00845

(22) International Filing Date: 7 May 2002 (07.05.2002)

(74) Agent: **CHOI, Kyu-Pal**; Halla Classic Building 4F,
824-11, Yeoksam-Dong, Kangnam-ku, Seoul 135-080
(KR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2001/25011 8 May 2001 (08.05.2001) KR

(81) Designated States (*national*): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): YON-
SUNG FINE CHEMICAL CO. [KR/KR]; 51-301,
Science Technology Park, Sungkyunkwan University,
300, Chonchon-Dong, Changan-Gu, Suwon, Changan-Gu,
Suwon, Kyunggi-Do 440-746 (KR).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HAM, Won-Hun**
[KR/KR]; Samik Apt. #12-1201, Cheongdam-Dong,
Kangnam-Gu, Seoul 135-100 (KR). **OH, Chang-Young**
[KR/KR]; 49-4, Seongsan-Dong, Mapo-Gu, Seoul
121-250 (KR). **KIM, Yong-Hyun** [KR/KR]; 73-73,
Hwagok 5-Dong, Kangseo-Gu, Seoul 157-015 (KR).
LEE, Yiu-Suk [KR/KR]; Woosung Apt. #304-407, 105,
Sadang-Dong, Dongjak-Gu, Seoul 156-090 (KR). **LEE,**

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 02/090324 A1


(54) Title: PROCESS FOR PREPARING PROSTAGLANDIN DERIVATIVES AND STEREOSPECIFIC STARTING MATERIAL THEREOF

(57) Abstract: The present invention relates to a process for effectively preparing prostaglandin derivatives and to a stereospecific alkyl halide containing 15S-alcohol group as a starting material.

PROCESS FOR PREPARING PROSTAGLANDIN DERIVATIVES AND STEREOSPECIFIC STARTING MATERIAL THEREOF

TECHNICAL FIELD

The present invention relates to a novel process for preparing a prostaglandin derivative of the following formula (1):




in which

R_1 represents H or C_1 - C_5 -alkyl,

X represents CH_2 , O, or S, and

R' represents C_2 - C_4 -alkyl; phenyl optionally substituted by halogen, C_1 - C_5 -alkyl, C_1 - C_4 -alkoxy, CF_3 , or C_1 - C_3 -aliphatic acylamino; 5- or 6-membered heterocycle containing one or more hetero atoms selected from a group consisting of nitrogen, oxygen and sulfur; C_3 - C_7 -cycloalkyl; or C_3 - C_7 -cycloalkenyl,

a 13,14-dihydro-PGF_{2a} ester derivative, and to an alkyl halide of the following formula (3a):



in which

Y' represents Br or I, and

R'' represents a hydroxy-protecting group, preferably trimethylsilyl, triethylsilyl, t-butyltrimethylsilyl, t-butylphenylsilyl, phenyldimethylsilyl, or tetrahydrofuranyl,

5 as a starting material.


BACKGROUND ART

10

Prostaglandin derivative of the above formula (1) is known in *Drugs of the Future*, 1992, 17(8), 691-704; *J. Med. Chem.*, 1993, 36, 243-248, etc. and a process of the following Reaction Scheme 1 published in WO 93/00329 can be mentioned as the typical synthesis thereof.

15

Reaction Scheme 1



In the above process, ω -chain is introduced into the starting material of Corey lactone by Wadworth-Emmons method, the ketone group at C15 position is reduced to an alcohol group, the remained double bond is reduced again using Pd, α -chain is introduced into a lactol as obtained through some consecutive reactions by Wittig reaction, and the terminal group is esterified.

However, the above process has been identified to have the following problems.

First, the diastereomeric mixture resulted from the introduction of ω -chain and reduction of the ketone group at C15 position comprises 15S-isomer and 15R-isomer in a ratio of 7:3. Therefore, stereoselectivity of the process is not satisfactory. Further, since these isomers can hardly be separated, the yield of the desired 15S-isomer is as very low as 38%.

Second, the yield of the esterification reaction of the terminal carboxylic acid after introduction of α -chain by Wittig reaction is reported to be less than 40%. This is the final step of the process, and so the low yield of the final step reaction may exert fatal and serious influence on the total efficacy of the synthesis.

Third, Corey lactone used as a starting material is very expensive, which makes the total process uneconomic.

25

DISCLOSURE OF INVENTION

Thus, the present inventors have extensively studied to develop an economic


and effective process for preparing the compound of formula (1) by solving the problems of prior arts as explained above. As a result, we newly designed an alkyl halide containing 15S-alcohol group and then completed a process for preparing the compound of formula (1), starting from 1,4-addition reaction at the α, β -unsaturated ketone group. According to the process of the present invention, the yield may be increased by using a stereospecific starting material instead of reducing the C15 ketone in a poor stereoselective manner; particularly, the low efficacy due to the low yield of esterification reaction of the terminal carboxylic acid may be improved; and there is no more need to use the expensive Corey lactone as a starting material. Therefore, the present invention provides an economic and effective synthetic process that is suitable for a mass production.

Therefore, an object of the present invention is to provide a process for preparing a prostaglandin derivative of formula (1).

It is another object of the present invention to provide an alkyl halide of formula (3a) as an effective starting material.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention relates to a process for preparing a compound of the following formula (1):




in which

R_1 represents H or C_1 - C_5 -alkyl,

5 X represents CH_2 , O, or S, and

R' represents C_2 - C_4 -alkyl; phenyl optionally substituted by halogen, C_1 - C_5 -alkyl, C_1 - C_4 -alkoxy, CF_3 , C_1 - C_3 -aliphatic acylamino; 5- or 6-membered heterocycle containing one or more hetero atoms selected from a group consisting of nitrogen, oxygen and sulfur; C_3 - C_7 -cycloalkyl; or C_3 - C_7 -cycloalkenyl, which comprises the first step wherein an alkyl

10 halide containing 15S-alcohol of the following formula (3):




in which

15 X and R' are defined as above,

Y represents a leaving group, preferably halogen, and

R'' represents a hydroxy-protecting group, preferably trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, phenyl dimethylsilyl, or tetrahydrofuranyl,


is converted into a cuprate thereof, and the cuprate compound is subjected to a
20 stereoselective 1,4-addition reaction to an α, β -unsaturated ketone compound of the following formula (2):



in which


R_1 is defined as above, and

- 5 R_2 represents a hydroxy-protecting group, preferably trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, phenyl dimethylsilyl, or tetrahydrofuranyl, to give a compound of the following formula (4):



10

in which X, R' , R'' , R_1 and R_2 are defined as above; the second step wherein the ketone group on the cyclopentanone ring of the compound of formula (4) is reduced using a metal hydride to give an α -alcohol compound of the following formula (5):



15


in which X, R' , R'' , R_1 and R_2 are defined as above; and the third step wherein the alcohol protecting groups on the cyclopentanone ring and ω -chain in the compound of

formula (5) are removed to give the compound of formula (1).

The process according to the present invention is depicted in the following Reaction Scheme 2 below.

5

Reaction Scheme 2



10

The process of the present invention proceeds through three steps, each of which is specifically explained in the following.

The first step : Preparation of the compound of formula (4) from the compounds of formulae (2) and (3)

15

The cuprate compound that is formed from the alkyl halide of formula (3) by

various methods as explained below is subjected to 1,4-addition reaction to the α, β - unsaturated ketone group of the compound of formula (2) to give the compound of formula (4). Here, the cuprate compound is introduced into the opposite side to the alkoxy group on the cyclopentenone due to the steric hindrance of the alkoxy group, resulting in a trans configuration with respect to the alkoxy group. The ω -chain thus introduced causes another steric hindrance to the α -chain of the cyclopentenone, whereby the compound of formula (4) in which α -chain and ω -chain have a trans configuration to each other is obtained.

10 The cuprate compound can be obtained from the compound of formula (3) by i) adding t-BuLi and then adding one substance selected from a group consisting of CuCN, (2-thienyl)Cu(CN)Li and MeCu(CN)Li to the compound of formula (3); or ii) adding one substance selected from a group consisting of CuBr · DMS, CuI and CuBr to a Grignard reagent that is formed from magnesium and the compound of formula (3).

15

Below, the process for preparing the compound of formula (4) from the alkyl halide of formula (3), where the cuprate compound acts as an intermediate, is specifically explained.

20

Method (a)

To 2 equivalents of the compound of formula (3) dissolved in a solvent is added 4 equivalents of t-BuLi at -78°C , which is then stirred. 1 equivalent of CuCN is added thereto. The temperature of the reaction solution is raised to -10°C to give a lower order-cuprate in a homogeneous state. This cuprate is cooled again to -78°C and then
25 reacted with 0.8-1.0 equivalent of the compound of formula (2).

Method (b)

To 1 equivalent of the compound of formula (3) dissolved in a solvent is added 2 equivalents of t-BuLi at -78°C , which is then stirred. 1 equivalent of (2-thienyl)Cu(CN)Li or MeCu(CN)Li is added thereto. The temperature of the reaction solution is slowly raised to -40°C over 30 minutes and then lowered again to -78°C to give a higher order-cuprate. This cuprate is reacted with 0.8-1.0 equivalent of the compound of formula (2).

Method ©


To 1 equivalent of the compound of formula (3) dissolved in a solvent is added 3-4 equivalents of Mg. The Grignard solution thus obtained is cooled to -78°C , and 0.3-1 equivalent of a substance selected from a group consisting of CuBr · DMS, CuI, and CuBr is added. The reaction mixture is stirred for 1 hour to give Normant reagent. This reagent is reacted with 0.8-1.0 equivalent of the compound of formula (2).

15

The compound of formula (3) used as a starting material in the above may be obtained by a process as exemplified in the following Reaction Scheme 3.

Reaction Scheme 3

20




That is, a nucleophile, preferably carbanion, phenol, thiophenol, or alkaline salt of phenol or thiophenol, is reacted with a chiral epoxide that can be prepared by a process known in *J. Chem. Research(S)*, 1983, 10-11 to open the ring in a stereospecific manner to give an alcohol compound having S stereochemistry. Particularly, in the case of carbanion, the alcohol compound can be obtained by adding 1~2 equivalents of a Grignard reagent and a catalytic amount (0.05~0.5 equivalent) of a substance selected from a group consisting of CuI, CuBr, and CuBr·DMS, stirring for 10 minutes, and slowly adding dropwise an epoxide in a solvent selected from a group consisting of diethylether, tetrahydrofuran and dimethylsulfide at reaction temperatures of -20~20°C.

Then, the alcohol is protected by a hydroxy-protecting group, preferably triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, phenyldimethylsilyl, or tetrahydrofuranyl, and the protecting group of the primary alcohol is removed. Particularly, the benzyl group used as a protecting group can be removed by a catalytic hydrogenation reaction in a solvent selected from a group consisting of methanol and ethanol, and in the presence of palladium as a catalyst; and the paramethoxybenzyl

group can be removed in a solvent mixture of dichloromethane and water in the presence of DDQ (2,3-dichloro-5,6-dicyano- 1,4-benzoquinone)(1.0 ~ 2.0 equivalent) at room temperature.

5 Finally, the primary alcohol is iodinated by I_2 , PPh_3 , and imidazole in a solvent mixture of acetonitrile and diethylether; or brominated by CBr_4 and PPh_3 , or reacting with methane sulfonyl chloride and triethylamine to give a methanesulfonate, which is then reacted with $LiBr$ in THF solvent.

10 Particularly, a compound of the following formula (3a) among the compound of formula (3):



15 in which Y' represents Br or I , and R'' represents a hydroxy-protecting group, is novel, and thus, it is another object of the present invention to provide the novel compound of formula (3a).

The compound of formula (2) can be prepared according to a process described
20 in *J. Org. Chem.*, **1978**, 43, 1641-1643, and (2-thienyl) $Cu(CN)Li$ or $MeCu(CN)Li$ can be obtained by adding the commercially available 2-thienyl lithium or $MeLi$ to $CuCN$.

In the above reaction, the solvent dissolving the compounds of formulae (2) and (3) is selected from a group consisting of tetrahydrofuran and diethylether, and the
25 solvent dissolving $t-BuLi$ is n -pentane. After the completion of each reaction, an

aqueous solution of $\text{NH}_4\text{Cl}/\text{NH}_4\text{OH}$ (9/1) is added to stop the reaction, Then, the reaction solution is extracted with diethylether, concentrated, and purified by column chromatography.

- 5 The second step : Preparation of the compound of formula (5) by reduction of the compound of formula (4)

 The compound of formula (4) obtained by the above 1,4-addition reaction has a ketone group, which can be reduced by various metal hydrides. As the metal hydrides that can be used, those having a heavy steric hindrance, preferably those selected from a group consisting of sodiumborohydride (NaBH_4), L-selectride, N-selectride and K-selectride, and particularly preferably L-selectride can be mentioned. This is because the hydride is apt to attack the cyclopentanone ring from the opposite direction with respect to the alkoxy group in the compound of formula (4) due to the steric hindrance of the alkoxy group, as the metal hydride is bulkier, to selectively give the desired α -alcohol. The reaction is carried out by adding 1~3 equivalents of L-selectride to 1 equivalent of the compound of formula (4) dissolved in a solvent at -78°C , stirring for 1 to 2 hours, and adding 30% H_2O_2 to stop the reaction. The reaction solution is stirred for 30 minutes at 0°C , extracted with diethylether, and concentrated to give the compound of formula (5) in a stereoselective manner. This compound is used in the next reaction without further purification. The solvent used in the above reaction is selected from a group consisting of tetrahydrofuran, diethylether and dichloromethane.

10
15
20

- The third step : Preparation of the compound of formula (1) through deprotection of the compound of formula (5)
- 25

 The two alcohol protecting groups in the compound of formula (5) can be removed under an acidic condition, or by using various fluorides (F) particularly when

the protecting group is silyl. More specifically, the following methods can be exemplified.

Method using an acidic condition

5

Method (a)

Deprotection is achieved by stirring for 24~48 hours at room temperature in the presence of excess NaHSO_4 in a solvent mixture of tetrahydrofuran and water (2:1).

10

Method using fluorides

Deprotection is achieved by

15

- stirring for 4 hours at 0°C ~room temperature in the presence of 2~4 equivalents of tetrabutylammonium fluoride ($\text{Bu}_4\text{N}^+\text{F}^-$) in a solvent of tetrahydrofuran (Method (b)); or

20

- reacting for 3~4 hours at 0°C with 2~10 equivalents of hydrogen fluoride pyridine (HF -pyridine) in a solvent of dichloromethane (Method (c)); or

- reacting for 4 hours with fluorosilicic acid (H_2SiF_6) in a solvent of acetonitrile according to DeShong method(*J. Org. Chem.*, 1992, 57, 2492) (Method (d)).

25

The present invention will be more specifically explained in the following examples. However, it should be understood that the following examples are intended to illustrate the present invention but not in any manner to limit the scope of the present invention.

Example 1 : Preparation of the compound of formula (3)**Preparation of 1-(4-methoxy-benzyloxy)-5-phenyl-pentan-3-ol**

5 Benzyl magnesium chloride (2.0M in THF) (64.6ml, 129.2mmol) was added to THF (77ml) and cooled to 0 °C. CuI (1.9g, 9.94mmol) was added, and the mixture was stirred for 10 minutes. 2-[2-(4-Methoxy-benzyloxy)-ethyl]-oxirane (19.5g, 99.4mmol) dissolved in THF (120ml) was slowly added dropwise thereto. The reaction mixture was stirred for 1 hour at room temperature and saturated aqueous NH₄Cl solution
10 (100ml) was added to separate the organic layer. The aqueous layer was extracted with diethylether (100ml× 2). The organic layers were combined, washed with saturated aqueous NaCl solution, dried over MgSO₄, filtered, and concentrated to give 1-(4-methoxy-benzyloxy)-5-phenyl-pentan-(S)-3-ol (30g) of a pale yellow oil. This compound was used in the next reaction without further purification.

15

Preparation of (S)-3-(t-butyldimethylsilyloxy)-5-(4-methoxy-benzyloxy)- 1-phenyl pentane

To DMF (390ml) were added 1-(4-methoxy-benzyloxy)-5-phenyl-pentan- (S)-3-ol (30g unpurified), imidazole (20.3g, 298.2mmol) and TBDMSCl (23g, 149.8mmol)
20 in the order, and the resulting mixture was stirred for 12 hours at room temperature. After completion of reaction, the reaction mixture was diluted with water (500ml) and extracted with ethyl acetate (100ml× 3). The organic layers were combined, washed with saturated aqueous NaCl solution, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel column chromatography (hexane:ethyl
25 acetate=20:1) to give (S)-3-(t-butyldimethylsilyloxy)-5-(4-methoxy- benzyloxy)-1-phenyl pentane (35g, Yield of 2 steps 85%) of a colorless and transparent oil.

Preparation of (S)-3-(t-butyldimethylsilyloxy)-5-phenyl-pentan-1-ol

(*S*)-3-(*t*-butyldimethylsilyloxy)-5-(4-methoxy-benzyloxy)-1-phenyl pentane (15g, 36.2mmol) was dissolved in a solvent mixture of dichloromethane-water (20:1) (252ml). DDQ (8.2g, 36.2mmol) was added thereto and the mixture was stirred for 30 minutes at room temperature. After completion of the reaction, saturated aqueous NaHCO₃ solution (100ml) was added to the reaction solution to stop the reaction. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The organic layers were combined, washed with saturated aqueous NaHCO₃ solution and saturated aqueous NaCl solution, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=15:1) to give (*S*)-3-(*t*-butyldimethylsilyloxy)-5-phenyl-pentan- 1-ol (10.6g, Yield 99%) of a yellow oil.

Preparation of (*S*)-3-(*t*-butyldimethylsilyloxy)-5-bromo-1-phenyl pentane

(*S*)-3-(*t*-butyldimethylsilyloxy)-5-phenyl-pentan-1-ol (10.6g, 36mmol) was dissolved in dichloromethane (80ml), triethylamine (12.6ml, 90.5mmol) was added, and the mixture was cooled to 0°C. Methane sulfonyl chloride (4.2ml, 54.3mmol) was slowly added dropwise and the resulting mixture was stirred for 12 hours. After completion of the reaction, the reaction mixture was diluted with water (50ml). The organic layer was separated and the aqueous layer was extracted with dichloromethane (50ml× 2). The organic layers were combined, washed with saturated aqueous NaCl solution, dried over MgSO₄, filtered, and concentrated. Without further purification, the residue was dissolved in THF (60ml). LiBr (7.86g, 90.5mmol) was added thereto and the mixture was refluxed for 4 hours. After completion of the reaction, water (80ml) was added to the reaction solution which was then extracted with diethylether. The organic layers were combined, washed with saturated aqueous NaCl solution, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=20:1) to give colorless and transparent (*S*)-3-(*t*-butyldimethylsilyloxy)-5-bromo-1-phenyl pentane (11g, Yield 86%).

Example 2 : Preparation of the compound of formula (4) according to Method (a)

5 (*S*)-3-(*t*-butyldimethylsilyloxy)-5-bromo-1-phenyl pentane (8mmol) was dissolved in diethylether (17ml) and cooled to -78 °C. *t*-BuLi (1.7M pentane solution, 16mmol) was slowly added dropwise thereto and the mixture was stirred for 20 minutes. Then, CuCN (4mmol) was added and the temperature of the reaction solution was slowly raised to -10 °C to make the solution homogeneous. The reaction solution was
10 cooled again to -78 °C. 7-[(*R*)-3-(*t*-butyldimethylsilyloxy)-5-oxo- cyclopent-1-enyl]-hept-5-enoic acid isopropyl ester (3.25mmol) dissolved in Et₂O (2.5ml) was slowly added dropwise and the mixture was stirred for 2 hours. The temperature of the reaction solution was raised to -30 °C, the reaction was stopped by the addition of 20ml of saturated aqueous NH₄Cl solution/28% aqueous NH₄OH solution (9/1), and the
15 reaction solution was warmed to room temperature. The reaction solution was diluted with diethylether, washed with water and brine, dried over Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=20:1) to give 11,15-O-bis(*t*-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGE_{2α} isopropyl ester in a yield of 82%.

20

Example 3 : Preparation of the compound of formula (4) according to Method (b)

25 (*S*)-3-(*t*-butyldimethylsilyloxy)-5-bromo-1-phenyl pentane (3.5mmol) was dissolved in diethylether (30ml) and cooled to -78 °C. *t*-BuLi (1.7M pentane solution, 7.35mmol) was added dropwise and the mixture was stirred for 10 minutes. (2-Thienyl)Cu(CN)Li (0.25M THF solution, 3mmol) which was newly prepared in another flask was added dropwise thereto. The reaction solution was slowly warmed to -40 °C

over 30 minutes and then cooled to -78°C . 7-[(*R*)-3-(*t*-butyldimethylsilyl oxy)-5-oxo-cyclopent-1-enyl]-hept-5-enoic acid isopropyl ester (2.7mmol) dissolved in diethylether(15ml) was slowly added dropwise thereto. Immediately after the dropwise addition, the reaction vessel was transferred to a cryostat of -45°C , and then
5 the reaction solution was warmed to -30°C over 30 minutes. After completion of the reaction, the reaction was stopped by adding 10ml of aqueous $\text{NH}_4\text{Cl}/\text{NH}_4\text{OH}(9/1)$ solution. The reaction solution was warmed to room temperature, diluted with diethylether, washed with water and brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum. The residue was purified by silica gel column chromatography
10 (hexane:ethyl acetate=20:1) to give 11,15-O-bis(*t*-butyldimethylsilyl) -13,14-dihydro-17-phenyl $\text{PGE}_{2\alpha}$ isopropyl ester in a yield of 83%.

Example 4 : Preparation of the compound of formula (4) according to Method ©

15

To a flask containing magnesium turning (25mmol) was introduced a solution wherein (*S*)-3-(*t*-butyldimethylsilyloxy)-5-bromo-1-phenyl pentane (8.3mmol) and 1,2-dibromoethane (0.05ml) were dissolved in THF (1.6ml). The reaction solution was warmed in a thermostat of 70°C to initiate formation of Grignard reagent, cooled again
20 to room temperature, and stirred for 30 minutes. This solution was diluted with THF (5ml), stirred for further 1.25 hour, and cooled to -78°C . $\text{CuBr}(\text{DMS})$ (2.0mmol) dissolved in dimethylsulfide (4ml) was added dropwise and the mixture was stirred for 1 hour. 7-[(*R*)-3-(*t*-butyldimethylsilyloxy)-5-oxo-cyclopent-1-enyl]- hept-5-enoic acid isopropyl ester (6.8mmol) dissolved in diethylether (7ml) was slowly added dropwise
25 thereto and the mixture was stirred for 1 hour. After completion of the reaction, the reaction solution was warmed to -30°C . The reaction was stopped by the addition of 20ml of aqueous $\text{NH}_4\text{Cl}/\text{NH}_4\text{OH}(9/1)$ solution, which was then warmed to room temperature. The reaction solution was diluted with diethylether, washed with water

and brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=20:1) to give 11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGE_{2a} isopropyl ester in a yield of 77%.

5

Example 5 : Preparation of the compound of formula (5)

11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGE_{2a} isopropyl ester (3.67mmol) was dissolved in THF (40ml), the reaction solution was cooled to -
10 78°C, and L-selectride (1M THF solution, 7.35mmol) was slowly added dropwise thereto. At this temperature the reaction solution was stirred for 2 hours and the reaction was stopped by the addition of hydrogen peroxide (30% aqueous solution, 16mmol). The reaction solution was stirred for 30 minutes at 0°C and extracted with diethylether. The organic layers were combined, washed with water and brine, dried over magnesium
15 sulfate, filtered, and concentrated under vacuum. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=10:1) to give 11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGF_{2a} isopropyl ester in the yield of 90%.

20 **Example 6 : Preparation of the compound of formula (1) according to Method (a)**

11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGF_{2a} isopropyl ester (4.32mmol) was dissolved in THF (25ml), $\text{NaHSO}_4 \cdot \text{H}_2\text{O}$ (17.3mmol) dissolved
25 in water (12.5ml) was added thereto, and the mixture was stirred for 24 hours at room temperature. After completion of the reaction, the reaction solution was extracted with water (20ml) and CH_2Cl_2 (85ml). The organic layers were combined, washed with saturated aqueous NaHCO_3 solution, dried over magnesium sulfate, filtered, and

concentrated. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=1:3) to give 13,14-dihydro-17-phenyl PGF_{2α} isopropyl ester in a yield of 75%.

5 **Example 7 : Preparation of the compound of formula (1) according to Method ③**

11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGF_{2α} isopropyl ester (1.22mmol) was dissolved in dichloromethane (25ml), HF·pyr (2.5ml) was
10 added thereto at 0°C, and the mixture was stirred for 3 hours. After completion of the reaction, the reaction solution was diluted with dichloromethane (10ml), which was then poured into saturated aqueous NaHCO₃ solution of 0°C. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The organic layers were combined, dried over magnesium sulfate, filtered, and concentrated. The
15 residue was purified by silica gel column chromatography (hexane:ethyl acetate=1:3) to give 13,14-dihydro-17-phenyl PGF_{2α} isopropyl ester in the yield of 82%.

Example 8 : Preparation of the compound of formula (1) according to Method ④

20

11,15-O-bis(t-butyldimethylsilyl)-13,14-dihydro-17-phenyl PGF_{2α} isopropyl ester (3.6mmol) was dissolved in acetonitrile (15ml), H₂SiF₆ (4ml, 25%wt aqueous solution) was added dropwise thereto in an ice bath, and the mixture was warmed to room temperature and stirred for 4 hours. After completion of the reaction, saturated
25 aqueous NaHCO₃ solution was added thereto. The aqueous layer was extracted with dichloromethane. The organic layers were combined, washed with water and brine, dried over magnesium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography (hexane:ethyl acetate=1:3) to give 13,14-dihydro-17-

phenyl PGF_{2α} isopropyl ester in the yield of 95%.

INDUSTRIAL APPLICABILITY


5

In the process of the present invention, a stereospecific starting material is used instead of reducing the ω -chain that may generate undesirable isomers and result in a yield decrease. Further, esterification of the carboxylic acid in the unstable
10 prostaglandin compound in the final step can be avoided, which in turn results in a yield increase in a considerable degree. That is, the total yield of the process according to the present invention is 38 to 51% based on the yields of examples above, which is much higher than the maximum yield of 15% of the previous process. Moreover, according to the present invention, the 15R-isomer resulted from the reduction of ω -
15 chain as an impurity, which can hardly be removed, is not generated, and also the production of impurity from the esterification reaction can be prevented. As a result, the prostaglandin derivative of formula (1) can be easily purified, and so can be synthesized in an economic and effective manner.

CLAIMS

1. A process for preparing a compound of the following formula (1):

5




in which

R₁ represents H or C₁-C₅-alkyl,

10 X represents CH₂, O, or S, and

R' represents C₂-C₄-alkyl; phenyl optionally substituted by halogen, C₁-C₅-alkyl, C₁-C₄-alkoxy, CF₃, C₁-C₃-aliphatic acylamino; 5- or 6-membered heterocycle containing one or more hetero atoms selected from a group consisting of nitrogen, oxygen and sulfur; C₃-C₇-cycloalkyl; or C₃-C₇-cycloalkenyl,

15 which comprises the first step wherein an alkyl halide containing 15S-alcohol of the following formula (3):



20

in which


X and R' are defined as above,

Y represents a leaving group, and

R'' represents a hydroxy-protecting group,

is converted into a cuprate thereof and the cuprate compound is subjected to a

stereoselective 1,4-addition reaction to an α, β -unsaturated ketone compound of the following formula (2):




in which

R_1 is defined as above, and


R_2 represents a hydroxy-protecting group,

10 to give a compound of the following formula (4):



in which X, R' , R'' , R_1 and R_2 are defined as above;

15 the second step wherein the ketone group on the cyclopentanone ring of the compound of formula (4) is reduced using a metal hydride to give an α -alcohol compound of the following formula (5):



in which X, R', R'', R₁ and R₂ are defined as above;

and the third step wherein the alcohol protecting groups on the cyclopentanone ring and ω-chain in the compound of formula (5) are removed to give the compound of formula (1).

2. The process of claim 1 wherein the compound of formula (3) is converted into its cuprate by


10 i) adding t-BuLi and then adding one substance selected from a group consisting of CuCN, (2-thienyl)Cu(CN)Li and MeCu(CN)Li to the compound of formula (3); or

ii) adding one substance selected from a group consisting of CuBr · DMS, CuI and CuBr to a Grignard reagent which is formed from magnesium and the compound of formula (3).

3. The process of claim 1 wherein the metal hydride is selected from a group consisting of sodiumborohydride (NaBH₄), L-selectride, N-selectride and K-selectride.

4. The process of claim 1 wherein the alcohol protecting groups are removed in the presence of H⁺ or F⁻.

5. A compound of the following formula (3a):



in which

5

Y' represents Br or I, and

R'' represents a hydroxy-protecting group.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR02/00845

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C07C 303/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean Patents and applications for inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Cas On line

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 364417, A1, (Stjerschants et al.), 18. 04. 90 see the whole document	1 ~ 4
A	WO 95/11003, A1, (Stjerschants et al.), 27. 04. 95 see the whole document	1 ~ 4

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 JULY 2002 (29.07.2002)

Date of mailing of the international search report

29 JULY 2002 (29.07.2002)

Name and mailing address of the ISA/KR



Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

KOH, Tae Wook

Telephone No. 82-42-481-5545

